

GTIN: 00363323314613

PHARMACY BULK PACKAGE –
NOT FOR DIRECT INFUSION

10 grams*
per Pharmacy Bulk Package

Vancomycin
Hydrochloride
for Injection, USP

Sample Label.
Please see package insert
for complete prescribing information.

NDC 63323-314-61

**Vancomycin
Hydrochloride**
for Injection, USP

10 grams*
per Pharmacy Bulk Package

PHARMACY BULK PACKAGE –
NOT FOR DIRECT INFUSION



**FRESENIUS
KABI**
Lake Zurich, IL 60047

www.fresenius-kabi.com/us

NDC 63323-314-61

**Vancomycin
Hydrochloride**
for Injection, USP

10 grams*
per Pharmacy Bulk Package

PHARMACY BULK PACKAGE –
NOT FOR DIRECT INFUSION

*Each bottle contains: Vancomycin hydrochloride
equivalent to 10 g vancomycin.

Usual adult dosage: 2 g daily, in divided doses.

STORE AT: 20° to 25°C (68° to 77°F) [see USP
Controlled Room Temperature].

After reconstitution: Use promptly. (Discard
bottle within 4 hours after entry.) See package
insert for full information.

The container closure is not made with natural
rubber latex.

**FRESENIUS
KABI**

NDC 63323-314-61

314061

**Vancomycin
Hydrochloride**
for Injection, USP

10 grams*
per Pharmacy Bulk Package

PHARMACY BULK PACKAGE –
NOT FOR DIRECT INFUSION

For intravenous infusion.

**MUST BE FURTHER DILUTED
AFTER RECONSTITUTION AND
BEFORE INFUSION. NOT TO BE
DISPENSED AS A UNIT.**

Preservative free.

Rx only

**FRESENIUS
KABI**

NDC 63323-314-61

**Vancomycin
Hydrochloride**
for Injection, USP

10 grams*
per Pharmacy Bulk Package

PHARMACY BULK PACKAGE –
NOT FOR DIRECT INFUSION

**IMPORTANT: READ INSERT FOR PRECAUTIONS
AND DIRECTIONS BEFORE USE.**

DIRECTIONS FOR USE:

Reconstitute with 95 mL of Sterile Water
for Injection using aseptic technique. Penetrate
one time after reconstitution using a sterile
dispensing set. **Each 5 mL of the resulting
solution contains 500 mg of vancomycin and
each 10 mL of the resulting solution contains
1 g of vancomycin. MUST BE FURTHER DILUTED
BEFORE USE.** This pharmacy bulk package is
for use in a Pharmacy Admixture Service.

Dispense aliquots from the bottle using a suitable
sterile dispensing set into infusion fluids under
a laminar flow hood using aseptic technique.
DISCARD BOTTLE NO LATER THAN 4 HOURS AFTER
INITIAL ENTRY (see package insert).

**FRESENIUS
KABI**

62526K

